



LONE OAK DENTAL
102 22nd Avenue S, Brookings, SD 57006
Ph: (605) 697-6212 | Fax: (605) 697-6232

Home Care After Scaling and Root Planing

These are some helpful suggestions that will increase your comfort and help you to receive the maximum benefit from periodontal treatment.

Tenderness is normal. To reduce tenderness and promote healing, every two to three hours, rinse with warm salt water: *½ teaspoon salt in a 4 ounce glass of water.*

Avoid brushing or flossing the treated area for 12 hours. However, after 12 hours it is very important that you continue to brush well. Please be careful brushing and use a toothbrush with soft bristles. You may moisten the brush with warm water if tissues are tender.

Tylenol or Ibuprofen may be used as recommended for discomfort. Doses of 600 mg of ibuprofen every 6 hours for the first 2-3 days should help with tender gums.

Highly nutritious food is necessary for the healing process. **Avoid foods that require excessive chewing, also, sticky, crunchy or coarse foods.**

Tooth sensitivity is normal and temporary. You may use a desensitizing toothpaste such as Sensodyne or any major brand for sensitivity relief.

You may also **gently massage** the areas treated with your washed fingers. This will increase circulation and promote healing.

Your **periodontal maintenance visits** are crucial to preventing your gum disease from progressing. Please keep your appointments with us at 3-4 month intervals.

Should you experience prolonged discomfort or have any questions, please contact us!